

# Risk aggregation: A general approach via the class of Generalized Gamma Convolutions

Justin Miles <sup>\*</sup>, Edward Furman<sup>\*,\*\*,1</sup>, and Alexey Kuznetsov<sup>\*</sup>

*<sup>\*</sup>Department of Mathematics and Statistics, York University,  
Toronto, Ontario M3J 1P3, Canada*

*<sup>\*\*</sup>School of Mathematical and Statistical Sciences, Western University,  
London, Ontario N6A 5B7, Canada*

**Abstract.** Risk aggregation is virtually everywhere in insurance applications. Indeed, in the vast majority of situations insurers are interested in the properties of the sums of the risks they are exposed to, rather than in the stand-alone risks per se. Unfortunately, the problem of formulating the probability distributions of the aforementioned sums is rather involved, and as a rule does not have an explicit solution. As a result, numerous methods to approximate the distributions of the sums have been proposed, with the moment matching approximations (MMAs) being arguably the most popular. The arsenal of existing MMAs is quite impressive and contains such very simple methods as the normal and shifted-gamma approximations that, respectively, match the first two and three moments, only, as well as such much more intricate methods as the one based on the mixed Erlang distributions [H.Cossette, D. Landriault, E. Marceau, and K. Moutanabbir (2016). Moment-based approximation with mixed Erlang distributions. *Variance* **10**(1), 166 – 182]. Note however that in practice the sums of insurance risks can have numerous and just a few summands; in the latter case the normal approximation is very questionable. Also, in practice the distributions of the stand alone risks can be light-tailed or heavy-tailed; in the latter case moments of higher orders (e.g.,  $\geq 2$ ) may not exist, and so the approximation based on mixed Erlang distributions is of limited usefulness.

In this paper we put forward a refined MMA method for approximating the distributions of the sums of insurance risks. Our method approximates the distributions of interest to any desired precision, works equally well for light and heavy-tailed distributions, and is reasonably fast irrespective of the number of the involved summands.

*Key words and phrases:* Generalized gamma convolution, Padé approximation, Gaver-Stehfest algorithm, individual risk model, collective risk model, policy modification

---

<sup>1</sup>Corresponding author. Email: [efurman@mathstat.yorku.ca](mailto:efurman@mathstat.yorku.ca)

# 1 Introduction

Risk aggregation is of fundamental importance for insurance. This is because risk aggregation is in fact a precursor of risk pooling, a principle that is seen by some as the insurance's reason d'être. To see how crucial risk aggregation is for risk pooling, consider a group of  $n \in \mathbf{N}$  individuals (also, business lines, risk components in a portfolio of risks, etc), where each one of  $i \in \{1, \dots, n\}$  faces a risk represented by the random variable (RV)  $X_i$ , then a sharing rule  $Y(X_1, \dots, X_n)$  is called risk pooling if it is a function of the aggregate risk  $X_1 + \dots + X_n$ , only, that is  $Y(X_1, \dots, X_n) = Y(X_1 + \dots + X_n)$  (e.g., [Aase, 2004](#), for details). Hence, it is clear that to reap the benefits of risk pooling, insurers must study and understand risk aggregation thoroughly. From now and on,  $X_1, \dots, X_n$  stand for insurance risk RVs, and  $X_1 + \dots + X_n := S_n$  denotes the associated aggregate risk.

Specific examples of risk aggregation are naturally abundant in all areas of insurance business. For classical applications, one has to go no further than the renowned individual (e.g., [Dhaene and Vincke, 2004](#)) and collective (e.g., [Goovaerts, 2004](#)) risk models. Specifically, in the case of the individual risk models (IRMs), for a fixed  $n$  as hitherto and independent but not identically distributed risk RVs  $X_1, \dots, X_n$ , the aggregate risk is as before denoted by  $S_n$ . Also, in the case of the collective risk models (CRMs), the number of risks, denoted by  $N$ , is assumed random and independent of identically distributed and mutually independent (IID) RVs  $X_1, \dots, X_N$ ; the aggregate risk is then denoted by  $S_N := X_1 + \dots + X_N$ .

In order to comprehend the implications of risk aggregation, and merely comply with the norms of the risk informed decision making, insurers are concerned with the stochastic properties of the RVs  $S_n$  and  $S_N$ , that is with the corresponding cumulative distribution functions (CDFs). Whether the IRM or CRM are considered, it is often tempting to approximate these CDFs with the use of the Lindeberg – Lévy central limit theorem (CLT). To this end, arguments like: (i) the number of risks is *large*, (ii) the risks are *not too* heterogeneous, and (iii) the distributions of the risk RVs are *not too* skewed are quite common. Unfortunately, none of the above must be true in reality. Furthermore, statements (ii) and (iii) are often violated as insurance risk RVs due to distinct risk sources can be very unlike, and, as a rule, positively skewed. Another problem with the aforementioned variant of the CLT is that there are situations (not rare) where the risk RVs of interest have infinite second moments (e.g., [Seal, 1980](#), and references therein). We refer to [Brockett \(1983\)](#) and references therein for some interesting examples of how the CLT is misused in insurance applications.

The standard CLT-based approximation of the aggregate risk's CDF, a.k.a. the normal approximation (NA), can be considered a moment matching approximation (MMA) that hinges on the first two moments only. A generalization of NA that incorporates skewness is the so-called normal-power (NP) approximation (e.g., [Ramsay, 1991](#)). An alternative approach to count for skewness that is of great popularity among practising actuaries is the shifted-gamma approximation (SGA), which aims to match

the first three moments (Hardy, 2004). Clearly, the choice of how many moments to match is somewhat ad-hoc. Thus more general approximations that aim at matching an arbitrary number of moments have been proposed (e.g., Cossette et al., 2016, for a recent reference).

Admittedly, MMAs, both the ones mentioned above and others alike, are convenient and intuitive to convey to upper management, yet, rather problematic. For instance, even the two moments-based NA method requires finite second moments, and it is inapplicable otherwise. The approach of Cossette et al. (2016) achieves better accuracy at the price of requiring the finiteness of higher order moments. In addition, in the latter case, the method is often rather computationally intensive.

In this paper, we put forward a new efficient method to approximate the CDFs of the aggregate risk RVs  $S_n$  and  $S_N$ . Our approach approximates the CDFs of interest to any desired precision, works equally well for light and heavy-tailed CDFs, and is reasonably fast irrespective of the number of summands. We organize the rest of this paper as follows: Section 2 provides a high-level overview of our approach, and Sections 3, 4, and 5 explain in detail its three main pillars, which are, respectively, the class of Padé approximations, the family of Generalized Gamma Convolutions, and the Gaver-Stehfest algorithm. The theory we propose is then elucidated by a variety of practical examples borrowed from Bahnemann (2015). More specifically, in Section 6 we demonstrate the effectiveness of the new MMA in the context of stand-alone risk RVs first, and then in the context of aggregate risk RVs with and without policy modifications.

## 2 Brief description of the method

In order to outline the essence of our proposed technique in the simplest possible manner, in this section we consider the framework of the IRM, however the ideas are equally applicable in the context of the CRM (Section 6 of this paper). Let  $X_1, \dots, X_n$  be positive and mutually independent RVs with arbitrary corresponding CDFs  $F_1, \dots, F_n$ . Also denote by  $\phi_i(z) := \mathbb{E}[\exp(-zX_i)]$  the Laplace transform (LT) of the RV  $X_i$ ,  $i = 1, \dots, n$ . Our goal is to approximate the CDF  $F$  of the aggregate RV  $S = X_1 + \dots + X_n$ .

Let  $\phi(z) := \mathbb{E}[\exp(-zS)]$  (also,  $\mathcal{L}(z)$ ) denote the LT of the aggregate risk RV  $S$ , then we readily have

$$\phi(z) = \prod_{i=1}^n \phi_i(z) \tag{1}$$

and

$$F(x) = \mathcal{L}^{-1} \left\{ \frac{\phi(z)}{z} \right\} (x), \quad x \geq 0. \tag{2}$$

Hence, by combining (1) and (2), we are able to obtain the desired approximation of the CDF  $F$  given that there exist reliable methods to: (i) approximate each LT  $\phi_i(z)$ , and (ii) invert LTs. Next in Sections 3 - 5, we demonstrate that (i) and (ii) can be achieved very successfully. Namely, in Section 3, we approximate

the Laplace transform  $\phi_i$  with the help of the Laplace transform of certain gamma convolutions, and we utilize the machinery of Padé approximations to determine the involved shape and rate parameters. In Section 4, we reintroduce the family of Generalized Gamma Convolutions, and, finally, in Section 5, we briefly explain the Gaver-Stehfest method to invert Laplace transforms. The proposed MMA may at the first glance seem complex, however, when implemented, it is utterly user-friendly and requires minimal human involvement.

### 3 Padé approximations

To start off, we note that (1) requires to approximate the LT of the RV  $X_i$  (the CDF of the RV  $X_i$  is then established via (2)). We accomplish this with the help of  $m$ -fold convolutions of Gamma-distributed RVs, succinctly  $\Gamma_i \sim \text{Gamma}(\alpha_i, \beta_i)$ ,  $i = 1, \dots, m$ . In other words, we seek to choose the parameters  $\{\alpha_i\}_{1 \leq i \leq m}$  and  $\{\beta_i\}_{1 \leq i \leq m}$ , so that the CDF of the approximant of order  $m$  RV  $\tilde{X}_{i,(m)} := \sum_{i=1}^m \Gamma_i$  is *close* in an appropriate sense to the CDF of the RV  $X_i$   $i = 1, \dots, n$ . In the rest of the paper, we omit the subscript  $(m)$  whenever the order of the approximation is fixed. Also, for the sake of the expositional simplicity of the discussion in the present section, and sometimes thereafter, we omit the subscript  $i$  and write  $X$  instead of  $X_i$ , and  $\phi$  instead of  $\phi_i$ .

In terms of LTs, we seek to choose the parameters  $\{\alpha_i\}_{1 \leq i \leq m}$  and  $\{\beta_i\}_{1 \leq i \leq m}$ , so that the approximant LT

$$\tilde{\phi}(z) := \mathbb{E} \left[ \exp \left( -z \sum_{i=1}^m \Gamma_i \right) \right] = \prod_{i=1}^m (1 + z/\beta_i)^{-\alpha_i}$$

is *close* to the function  $\phi(z) = \mathbb{E}[\exp(-zX)]$ . Our method is essentially an MMA. Note that the moments of  $X$  can be computed as  $\mathbb{E}[X^k] = (-1)^k \phi^{(k)}(0)$ , thus matching the first  $m$  moments of  $\tilde{X}$  and  $X$  is equivalent to matching the derivatives (of order  $k = 1, 2, \dots, m$ ) of  $\tilde{\phi}$  and  $\phi$  at  $z = 0$ . However, our approach allows for a number of important improvements.

**Note 1.** *A significant innovation of our method is that we match the derivatives of  $\tilde{\phi}$  and  $\phi$  not at  $z = 0$  but at some point  $z = z^* > 0$ . This is crucial if we want our technique to be applicable to risk RVs with heavy tails, for which the moments (and thus the derivatives  $\phi^{(k)}(0)$ ) may fail to exist. If the risk RV  $X$  has exponentially light tails, we may as well choose  $z^* = 0$ , but in general  $z^*$  must be strictly positive.*

**Note 2.** *Another distinguishing feature of our approach is that  $\tilde{\phi}$  converges to  $\phi$  uniformly and exponentially fast, and in particular the approximant RV  $\tilde{X}$  converges in distribution to the RV  $X$ .*

**Note 3.** *Our method requires  $2m$  parameters (unique up to permutation) to match the first  $2m$  moments of the RVs  $\tilde{X}$  and  $X$ . As at least  $2m$  parameters are required to complete this task, the method we put forward herein is optimal in this sense.*

### 3.1 Approximation of order two

In order to explain the main ideas behind our algorithm, let us consider a simple case where  $m = 2$  and  $z^* = 1$ . In this case the problem reduces to the following one: we want to find four positive numbers  $\alpha_1$ ,  $\alpha_2$ ,  $\beta_1$  and  $\beta_2$  such that the derivatives of the LT  $\phi$  at  $z = 1$  coincide with derivatives of the approximant LT

$$\tilde{\phi}_{(2)}(z) = (1 + z/\beta_1)^{-\alpha_1}(1 + z/\beta_2)^{-\alpha_2}$$

also at point  $z = 1$ . In order to compute the four constants  $\alpha_1$ ,  $\alpha_2$ ,  $\beta_1$  and  $\beta_2$  we need to have at least four equations, so now we need to solve the following system

$$\frac{d^k}{dz^k}(1 + z/\beta_1)^{-\alpha_1}(1 + z/\beta_2)^{-\alpha_2}\Big|_{z=1} = \phi^{(k)}(1), \quad k \in \{1, 2, 3, 4\}. \quad (3)$$

If we were to compute the derivatives in the left-hand side of (3) and then to simplify the resulting equations, we would have obtained a fairly complicated system of four nonlinear equations in  $\alpha_1$ ,  $\alpha_2$ ,  $\beta_1$  and  $\beta_2$ . However, analysing these equations theoretically would not be feasible due to their complexity and even solving them numerically would be a major problem.

In order to avoid the complexity, we take logarithms before differentiating. Thus, instead of matching the derivatives  $\phi^{(k)}$  and  $\tilde{\phi}_{(2)}^{(k)}$  for  $k = 1, 2, 3, 4$ , we match the derivatives of  $\ln(\phi(z))$  and  $\ln(\tilde{\phi}_{(2)}(z))$  of order  $k = 1, 2, 3, 4$ . The end result is clearly the same, but computing  $\alpha_1$ ,  $\alpha_2$ ,  $\beta_1$  and  $\beta_2$  is much simpler, as we demonstrate in a moment. With this change, the system of four equations for finding  $\alpha_1$ ,  $\alpha_2$ ,  $\beta_1$  and  $\beta_2$  looks as follows:

$$\frac{d^k}{dz^k} \ln \left[ (1 + z/\beta_1)^{-\alpha_1}(1 + z/\beta_2)^{-\alpha_2} \right] \Big|_{z=1} = \frac{d^k}{dz^k} \ln[\phi(z)] \Big|_{z=1}, \quad k \in \{1, 2, 3, 4\}. \quad (4)$$

Let us denote  $s_k := -(1/k!) \frac{d^{k+1}}{dz^{k+1}} \ln[\phi(z)] \Big|_{z=1}$ . As we see later, it is easy to compute  $s_k$  numerically in each case of interest, thus for now we treat these numbers as known quantities. Computing the derivatives in the left-hand side of (4), we obtain a system of four equations

$$\frac{\alpha_1}{(1 + \beta_1)^k} + \frac{\alpha_2}{(1 + \beta_2)^k} + \frac{\alpha_2}{(1 + \beta_2)^k} + \frac{\alpha_2}{(1 + \beta_2)^k} = (-1)^k s_{k-1}, \quad k \in \{1, 2, 3, 4\}. \quad (5)$$

At this stage, that is in order to solve the nonlinear equations in (5) and find  $\alpha_1$ ,  $\alpha_2$ ,  $\beta_1$  and  $\beta_2$ , we employ the toolbox of *Padé approximations*.

Namely, we introduce yet another function  $\psi(z) := -\phi'(z)/\phi(z)$ . Then system (4) is equivalent to

$$\frac{d^k}{dz^k} \left[ \frac{\alpha_1}{z + \beta_1} + \frac{\alpha_2}{z + \beta_2} \right] \Big|_{z=1} = \psi^{(k)}(1), \quad k \in \{0, 1, 2, 3\}. \quad (6)$$

Note that we can write

$$\frac{\alpha_1}{z + \beta_1} + \frac{\alpha_2}{z + \beta_2} = \frac{A + B(z - 1)}{1 + C(z - 1) + D(z - 1)^2} \quad (7)$$

for some constants  $A$ ,  $B$ ,  $C$  and  $D$ . Now we can express system of four equations (6) in an equivalent way by saying that the first four terms of the Maclaurin expansion of the rational function

$$\frac{P(w)}{Q(w)} := \frac{A + Bw}{1 + Cw + Dw^2}$$

must match the corresponding terms of the Maclaurin expansion

$$\psi(1 + w) = s_0 + s_1w + s_2w^2 + s_3w^3 + \dots,$$

in other words we have a single equation

$$\frac{A + Bw}{1 + Cw + Dw^2} = s_0 + s_1w + s_2w^2 + s_3w^3 + O(w^4), \quad w \rightarrow 0, \quad (8)$$

(note that we have introduced here a new variable  $w := z - 1$ ). Equation (8) tells us that the rational function  $P(w)/Q(w)$  is a  $[1/2]$  Padé approximation to the function  $\psi(1 + w)$ . In general, a  $[p/q]$  Padé approximation to a function  $f$  is a rational function  $P(w)/Q(w)$  (with  $\deg(P) = p$  and  $\deg(Q) = q$ ) that has the same first  $p + q + 1$  terms in Maclaurin expansion as the function  $f(w)$ .

It turns out that single equation (8) contains all the information necessary for finding the constants  $A$ ,  $B$ ,  $C$  and  $D$ . By multiplying both sides of equation (8) by  $Q(w) = 1 + Cw + Dw^2$  we obtain

$$A + Bw = (1 + Cw + Dw^2)(s_0 + s_1w + s_2w^2 + s_3w^3 + O(w^4)), \quad w \rightarrow 0. \quad (9)$$

Identifying the coefficients in front of powers of  $w$  in both sides of the above equation we obtain a system of four linear (!) equations

$$\begin{cases} A = s_0, \\ B = Cs_0 + s_1, \\ 0 = Ds_0 + Cs_1 + s_2, \\ 0 = Ds_1 + Cs_2 + s_3. \end{cases} \quad (10)$$

Note that this system of four linear equations can be solved very easily by first finding  $C$  and  $D$  from the third and fourth equations, and then substituting these results into the first and second equations would give us the values of  $A$  and  $B$ .

Now that we have found  $A$ ,  $B$ ,  $C$ , and  $D$ , and so we can compute  $\beta_1$  and  $\beta_2$  by noting that

$$(1 + \beta_1 + w)(1 + \beta_2 + w) = 1 + Cw + Dw^2,$$

and then the constants  $\alpha_1$  and  $\alpha_2$  can be found by the formula  $\alpha_i = P'(-1 - \beta_i)$ . Remarkably, all we have done so far is the partial fraction decomposition for the rational function  $P(w)/Q(w)$  as in (7). This process, which can be done *by hand*, has provided us with the desired constants  $\alpha_1$ ,  $\alpha_2$ ,  $\beta_1$  and  $\beta_2$ , and so with the approximant LT  $\tilde{\phi}_{(2)}$ .

### 3.2 Extension to the approximation of any order

To generalize the method described in the previous subsection to arbitrary orders  $m$  and arbitrary choice  $z^*$ , we follow the same steps and arrive at the problem of finding  $[m - 1/m]$  Padé approximant to the function  $\psi(z^* + w)$ , that is, instead of (7) and (8) we have

$$\begin{aligned} \sum_{i=1}^m \frac{\alpha_i}{z^* + \beta_i + w} &= \frac{a_0 + a_1 w + \dots + a_{m-1} w^{m-1}}{1 + b_1 w + b_2 w^2 + \dots + b_m w^m} \\ &= s_0 + s_1 w + s_2 w^2 + \dots + s_{2m-1} w^{2m-1} + O(w^{2m}), \end{aligned} \quad (11)$$

where  $s_k := (1/k!) \psi^{(k)}(z^*)$ . This allows us to find the coefficients  $b_i, a_i$  by solving a system of linear equations similar to (10) and then to obtain the required numbers  $\{\alpha_i\}_{1 \leq i \leq m}$  and  $\{\beta_i\}_{1 \leq i \leq m}$  by doing the partial fraction decomposition in (11).

The main input for this algorithm is the sequence of coefficients  $s_k = (1/k!) \psi^{(k)}(z^*)$ . These coefficients are typically not available in closed form, but they can be easily computed numerically. Indeed, we first compute the numbers

$$g_k := \phi^{(k)}(z^*) = (-1)^k \int_0^\infty x^k e^{-z^* x} dF(x), \quad k \geq 0, \quad (12)$$

to a high precision by a numerical quadrature (the double-exponential quadrature of [Takahasi and Mori \(1974\)](#) is particularly well-suited for such calculations). Next we observe that

$$\frac{d}{dz} \phi(z) = -\psi(z) \phi(z). \quad (13)$$

Rewriting this identity in terms of Taylor series centred at  $z^*$  gives us

$$\sum_{k \geq 0} g_{k+1} (z - z^*)^k / k! = - \left( \sum_{n \geq 0} s_n (z - z^*)^n \right) \times \left( \sum_{k \geq 0} g_k (z - z^*)^k / k! \right).$$

Comparing the constant term in the Taylor series in the left-hand side and the right-hand side of the above equation we find that  $s_0 = -g_1/g_0$ , and comparing the coefficients in front of  $(z - z^*)^k$  gives us

$$s_k = -\frac{1}{g_0} \left( \frac{g_{k+1}}{k!} + \sum_{i=0}^{k-1} s_i \frac{g_{k-i}}{(k-i)!} \right), \quad (14)$$

which allows to compute  $s_k$  recursively for all  $k \geq 1$ .

### 3.3 A simple numerical example and a question arising from it

To illustrate our method on a numerical example of actuarial interest, let us assume that  $X$  is distributed Weibull with the probability density function (PDF)

$$f(x) = (3/4)x^{-1/4} \exp(-x^{3/4}), \quad x \geq 0.$$

Weibull distribution has been frequently chosen to model insurance risks ([Mikosch, 2009](#); [Klugman et al., 2012](#), for general discussions).

We fix  $m = 2$  and  $z^* = 1$ ; by computing numerically the integral in (12) we calculate

$$g_0 \approx 0.5193711 \quad g_1 \approx -0.2123717, \quad g_2 \approx 0.2179689, \quad g_3 \approx -0.3665409, \quad g_4 \approx 0.8649004.$$

Then we use (14) to find

$$s_1 \approx 0.4089017, \quad s_2 \approx -0.0868391, \quad s_3 \approx 0.0630061, \quad s_4 \approx -0.0478247$$

and solving system of linear equations (10) we obtain

$$A \approx 0.4089017, \quad B \approx -0.2292057, \quad C \approx 0.7729116, \quad D \approx 0.0100583.$$

The polynomial  $Q(w) = 1 + Cw + Dw^2$  has two roots  $\approx -75.526356$  and  $\approx -1.3163588$ , thus we find  $\beta_1 \approx 74.526356$  and  $\beta_2 \approx 0.3163588$ . Using the formula  $\alpha_i = P'(-1 - \beta_i)$  we find  $\alpha_1 \approx 22.6439903$  and  $\alpha_2 \approx 0.1435963$ . Thus we have found an approximation to the Weibull distribution of interest. Namely, we have approximated  $X$  with the help of  $\tilde{X}_{(2)}$ , such that the latter RV is equal in distribution to  $\Gamma_1 + \Gamma_2$ , with  $\Gamma_i \sim \text{Gamma}(\alpha_i, \beta_i)$ ,  $i = 1, 2$ .

We conclude this section with an important observation, which paves the path to the introduction of the class of GGCs latter on in the next section. Namely, let the RV  $X$  be distributed Weibull with the following PDF (the shape parameter is now greater than one)

$$f(x) = (3/2)x^{1/2} \exp(-x^{3/2}), \quad x \geq 0.$$

For this set-up, we find that  $\alpha_1 \approx -29.421049$  and  $\beta_1 \approx -66.009893$ , which shows that our algorithm does not always give a legitimate approximation in the form  $\Gamma_1 + \Gamma_2$ , with  $\Gamma_i \sim \text{Gamma}(\alpha_i, \beta_i)$ . This outcome is rather unfortunate, but similar inconveniences have been encountered in the context of other MMAs (e.g. [Cossette et al., 2016](#)).

This raises an important question:

Does there exist a (rich) class of CDFs for which the Padé approximations described above always yield meaningful results?

In the next section we show that the answer to this question is in affirmative, and we also discuss the convergence of our refined MMA method.

## 4 Generalized Gamma Convolutions

The class of Generalized Gamma Convolutions comprises distributions which are weak limits of the RVs of the form  $\sum_{i=1}^m \Gamma_i$ , where  $\Gamma_i \sim \text{Gamma}(\alpha_i, \beta_i)$  and  $\alpha_i > 0$ ,  $\beta_i > 0$ ,  $i = 1, \dots, m$ . It seems that



GGCs were first used by Thorin in 1977, who employed their properties to prove that the log-normal distribution is infinitely divisible (e.g., [Bondesson, 1992](#), for an excellent discussion).

The most convenient way to describe the class of GGCs is via LTs. Note that for a RV  $\sum_{i=1}^m \Gamma_i := \tilde{X}_{(m)}$ , which is our approximant from Section 3, we have

$$\mathbb{E}[\exp(-z\tilde{X}_{(m)})] = \prod_{i=1}^m \mathbb{E}[\exp(-z\Gamma_i)] = \prod_{i=1}^m (1 + z/\beta_i)^{-\alpha_i} = \exp\left(-\int_0^\infty \ln(1 + z/t)U(dt)\right),$$

where  $U(dt)$  is a discrete measure having support at points  $\beta_i$  with mass  $U(\{\beta_i\}) = \alpha_i$ . Thus, the following result is not surprising.

**Proposition 1** ([Thorin \(1977\)](#), see also [Bondesson \(1992\)](#)). *The distribution on  $[0, \infty)$  of the r.v.  $X$  is a GGC if and only if its Laplace transform is*

$$\phi(z) := \mathbb{E}[\exp(-zX)] = \exp\left(-az - \int_0^\infty \ln(1 + z/t)U(dt)\right) \text{ for } \operatorname{Re}(z) \geq 0, \quad (15)$$

where  $a \in [0, \infty)$  is a constant, and  $U(dt)$  is a positive Radon measure, also called Thorin measure, which must satisfy

$$\int_0^\infty \min(|\ln(t)|, 1/t)U(dt) < \infty.$$

It turns out that the algorithm outlined in Section 3 always produces meaningful results when the RV being approximated has a CDF that belongs to the class of GGCs. By meaningful results we mean that the algorithm produces a set of *positive* numbers  $\alpha_i$  and  $\beta_i$  that determine the approximant  $\tilde{X}_{(m)} := \sum_{i=1}^m \Gamma_i$ , where  $\Gamma_i \sim \text{Gamma}(\alpha_i, \beta_i)$ . Moreover, it can be shown ([Furman et al., 2018](#)) that, as  $m \rightarrow +\infty$ , the LTs  $\mathbb{E}[\exp(-z\tilde{X}_{(m)})]$  converge to the LT  $\mathbb{E}[\exp(-zX)]$  exponentially fast and uniformly in  $z$  on compact subsets of  $\mathbb{C} \setminus (-\infty, 0]$ . One may wonder at this point whether the class of GGCs is rich enough. We note in this respect that it comprises, e.g., such important distributions for actuarial applications as gamma, inverse gamma, inverse Gaussian, Pareto, log-normal, and Weibull with the shape parameter less than one, among a great variety of other distributions.

To summarize the findings of Sections 3 and 4, we emphasize that our MMA (i) converges very fast, (ii) always provides legitimate outcomes if the risk RV to be approximated has a CDF in the class of GGCs, (iii) yields unique parameters of the approximant CDF, and (iv) is optimal in the sense that only  $2m$  parameters are required to match the first  $2m$  moments.

We conclude this section with outlining the cause for the algorithm described in Section 3 to be so well-suited for the RVs having CDFs in the class of GGCs. This reason in fact stems from the fact that the function  $\psi(z) = -d/dz \ln(\mathbb{E}[\exp(-zX)])$  is given by

$$\psi(z) = \int_0^\infty \frac{U(dt)}{t + z}, \quad (16)$$

which can be easily obtained from (15). Functions of the form  $\int_0^\infty (t+z)^{-1}U(dt)$  are called *Stieltjes functions*, and they have been shown to enjoy many nice analytical properties. In particular, it has been proved that Padé approximations to such functions always exist and converge exponentially fast and uniformly in  $z$  on compact subsets of  $\mathbb{C} \setminus (-\infty, 0]$  (see Baker and Graves-Morris (1996)). Since the algorithm outlined in Section 3 is essentially a Padé approximation method, this explains why RVs with CDFs in the class of GGCs fit perfectly within our approximation scheme.

## 5 Gaver-Stehfest algorithm

At this point of the discussion, we have hopefully convinced the reader that there is a mathematically sound way to approximate the risk RV  $X$  having a CDF in the class of GGCs with the help of an approximant RV  $\tilde{X}$ . The solution, however attractive, involves the corresponding LTs, that is  $\tilde{\phi} \approx \phi(z)$ . However, our ultimate goal is to find the CDF of the approximant RV  $\tilde{X}$ , denoted by  $\tilde{F}$ . Thus, the remaining step is to rediscover this CDF from the LT  $\tilde{\phi}$ . Doing that is a classical problem in analysis, and a variety of solutions exist. One popular method is to use Bromwich integral, which requires integration over a path in the complex plane. Another method, and this is what we do in the present paper, is via the Gaver-Stehfest algorithm. The main difference between the Gaver-Stehfest algorithm and the one based on the Bromwich integral, as well as most other Laplace inversion methods, is that it uses only the values of the Laplace transform on the positive real line and does not require any complex numbers. This method was invented in 1970 by Stehfest (Stehfest, 1970), by improving upon the earlier method of Gaver, and since then it has been successfully used in many areas of Applied Mathematics, including in Probability and Statistics (Abate and Whitt, 1992; Kou and Wang, 2003), Actuarial Science (Badescu et al., 2005) and Mathematical Finance (Schoutens and Damme, 2011).

The Gaver-Stehfest algorithm is very simple and easy to implement. To introduce it, consider a function  $f$  and its Laplace transform

$$\phi(z) := \int_0^\infty e^{-zx} f(x) dx,$$

where we assume that  $\phi(z)$  is finite for all  $z > 0$ . For all integers  $m \geq 1$  we define

$$f_m(x) := \ln(2)x^{-1} \sum_{k=1}^{2m} a_k(n) \phi(k \ln(2)x^{-1}), \quad x > 0, \quad (17)$$

where the coefficients are defined as follows:

$$a_k(m) := \frac{(-1)^{m+k}}{m!} \sum_{j=\lceil (k+1)/2 \rceil}^{\min(k,m)} j^{m+1} \binom{n}{j} \binom{2j}{j} \binom{j}{k-j}, \quad m \geq 1, 1 \leq k \leq 2m.$$

It is known (Kuznetsov, 2013) that the approximations  $f_m(x)$  converge to  $f(x)$  if  $f$  is continuous at  $x$  and of bounded variation in a neighbourhood of  $x$ . There is also a lot of numerical evidence that the approximations  $f_m(x)$  converge to  $f(x)$  very fast, provided that  $f$  is smooth enough at  $x$  (e.g., Abate and Whitt, 1992; Davies and Martin, 1979). When using the Gaver-Stehfest algorithm one should be careful with the loss of significant digits in the sum (17). This is due to the fact that the coefficients  $a_k(m)$  are very large (for large  $k$  and  $m$ ) and of alternating signs. This problem is readily solved by using any high-precision arithmetic package.

## 6 Illustrative examples with log-normal, Pareto and Weibull risks of varying tail thickness

In this section, we demonstrate the usefulness of our new MMA method by applying it to a number of examples of actuarial interest. More specifically, motivated by numerical examples in Bahnemann (2015), we consider applications to risks that are distributed log-normally, Pareto of the second kind, a.k.a., Lomax, and Weibull. These distributions have been routinely chosen to model insurance risks (e.g., Kleiber and Kotz, 2003; Mikosch, 2009; Klugman et al., 2012, for a general discussion).

In what follows, we denote the CDFs of the log-normal, Lomax, and Weibull distributions, by, respectively,  $F_L$ ,  $F_P$ , and  $F_W$ . All these CDFs have explicit forms given by

$$F_L(x; \mu, \sigma) = \frac{1}{2} \left[ 1 + \operatorname{erf} \left( \frac{\ln x - \mu}{\sigma \sqrt{2}} \right) \right], \quad x > 0, \mu \in \mathbb{R}, \sigma > 0, \quad (18)$$

$$F_P(x; \alpha, \beta) = 1 - \left( \frac{\beta}{x + \beta} \right)^\alpha, \quad x \geq 0, \alpha > 0, \beta > 0, \quad (19)$$

$$F_W(x; \beta, \delta) = 1 - e^{-(x/\beta)^\delta}, \quad x \geq 0, \beta > 0, \delta > 0. \quad (20)$$

In the above,  $\sigma$ ,  $\alpha$  and  $\delta$  are shape parameters that stipulate how heavy-tailed the corresponding distributions are. Namely, in the case of the log-normal distribution, larger values of  $\sigma$  imply heavier right tails, whereas in the case of the Lomax and Weibull distributions, smaller values of  $\alpha$  and  $\delta$ , respectively, suggest heavier right tails. Interestingly, there has been ample practical evidence that these are the parametrizations that correspond to the heavier tails that are of particular interest in non-life insurance applications, yet in the theoretical literature the light-tailed examples prevail (e.g., Cossette et al., 2016; Tao et al., 2016, for recent references). In the rest of this section, we consider both heavy-tailed and light-tailed parametrizations of the CDFs  $F_L$ ,  $F_P$ , and  $F_W$ . Our choices of parameters are inspired by Bahnemann (2015).

## 6.1 Stand-alone risks

We begin by approximating CDFs (18), (19), and (20). Genuinely speaking, these approximations are not the ultimate goal of this paper, and hence we remind the reader that: (i) after we have the approximations for the just-mentioned CDFs of the stand-alone risks RVs, the CDFs of the aggregate risk RVs are obtained with the help of equations (1) and (2), (ii) in the context of the stand-alone risks, we are able to evaluate the accuracy of an approximation by comparing the approximant CDF with the actual CDF, and thus the discussion in this subsection can serve as an important evaluation of the various MMAs tested here.

To explore the effectiveness of the distinct MMAs, we employ the Kolmogorov-Smirnov (KS) metric to measuring how close the distribution of an approximation is to the desired distribution. The KS distance ( $d_{KS}$ ) of an approximant RV  $\tilde{X}$  (with CDF  $\tilde{F}$ ) to the RV  $X$  (with CDF  $F$ ), denoted  $d_{KS}(\tilde{F}, F)$ , is given by

$$d_{KS}(\tilde{X}, X) = \sup_{x \geq 0} \left| \tilde{F}(x) - F(x) \right|.$$

This metric yields the worst distance and, thus, small values of  $d_{KS}$  suggest the approximation is good on the entire domain. However, on a different note, relatively large values of  $d_{KS}$  do not necessarily mean the approximation is worthless, as it can turn out very reasonable in some regions of the CDFs domain.

In the examples below, we illustrate the effectiveness of our approximation by comparing it with three simple MMAs: the normal approximation, the normal power approximation (NPA), and the shifted gamma approximation, when applicable. These three MMAs are referred to as ‘the most commonly cited in practice’ by Hardy (2004). In addition, we compare our approximation with the mixed Erlang distribution (MED) approach of Cossette et al. (2016), when applicable. The MED method uses the first  $m$  moments of a risk RV to determine a mixture of Erlang distributions for approximating this RVs CDF. We use the results in Cossette et al. (2016) for the log-normal distribution ( $\mu = 0$ ,  $\sigma = 0.5$ ) and apply their algorithm to our additional cases of interest (Pareto and Weibull), when applicable. (When implementing the MED method, we only considered moment-matching of orders  $m = 3$  and  $m = 4$ , as with  $m \geq 5$  the method requires a large number of computations.)

In what follows, we denote the CDF of our approximant RV  $\tilde{X}_{(m)}$  by  $\tilde{F}_{(m)}$ , we denote the mixed Erlang CDF by  $\tilde{F}_{W_{m,l}}$ , and the CDFs of the NA, NPA, and SGA methods by  $\tilde{F}_{NA}$ ,  $\tilde{F}_{NP}$ , and  $\tilde{F}_{SG}$ , respectively.

**Example 1.** To start off, let us consider log-normally distributed risk RVs. The log-normal distribution has been found appropriate for modelling losses originating from a great variety of non-life insurance risks (e.g., Mikosch, 2009; Klugman et al., 2012). More specifically, Kleiber and Kotz (2003) mention applications in property, fire, hurricane, and motor insurances, to name just a few (also, e.g., Dropkin, 1964; Bickerstaff, 1972; O’Neill and Wells, 1972). Furthermore the standard formula of the European Insurance and Occupational Pensions Authority explicitly assumes the log-normality of insurers’ losses

(EIOPA-14-322, 2014).

Table 1 provides the KS distances of several approximations for the parameter sets  $(\mu = 0, \sigma = 0.5)$ ,  $(\mu = 1.5240, \sigma = 1.2018)$ , and  $(\mu = 5.9809, \sigma = 1.8)$ . The choice of the larger value of the  $\sigma$  parameters is motivated by Example 5.8 in Bahnemann (2015), whereas the choice of the smaller one is due to Cossette et al. (2016); Dufresne (2007). As the parameter  $\sigma$  gets larger, it becomes increasingly difficult to obtain a valid approximation with any of the methods (this is perhaps the reason why in the examples manifesting in the academic literature, smaller values of the  $\sigma$  parameter are very common). With  $\sigma = 1.2018$  and  $\sigma = 1.8$ , the MED method does not produce potential solutions using the parameters  $m = 4$  and  $l = 70$ , in the former case, and  $m = 3$ , or 4 and  $l = 70$ , in the latter case. As suggested in Cossette et al. (2016), this can be remedied by increasing the parameter  $l$ ; however, increasing this parameter too much results in the method being computationally infeasible.

Figure 1 provides the plots of the approximating CDFs in Table 1 as well as the actual CDF  $F_L$  (in blue). Figures 1 a), c), and e) depict our approximations (dark green and light green, where light green corresponds to the better approximation) and the MED approximation (in red, when applicable). Figures 1 b), d), and f) depict the simple moment-matching methods: NA (orange), NPA (salmon), and SGA (black). Apparently, the NA and NPA approximations are inadequate for the more skewed log-normal cases, that is for the parametrizations  $(\mu = 1.5240, \sigma = 1.2018)$  and  $(\mu = 5.9809, \sigma = 1.8)$ . This is not surprising, as it is well-known that the NPA provides fairly accurate results for the skewness values not exceeding one (e.g. Ramsay, 1991), whereas the two aforementioned cases of the log-normal distribution lead to the skewness values of 11.23 and 136.38, respectively. In the context of the SGA, higher skewness values imply very small scale parameters, and these in turn result in a nearly vertical rise in the corresponding CDF, thus aggravating the accuracy significantly.

**Example 2.** The next distribution we consider is Lomax. As in the case of the log-normal distribution discussed earlier, there is ample of evidence that the Lomax distribution is an adequate model to describing non-life insurance risks. We refer to Seal (1980) for a list of references, and in particular to Benckert and Sternberg (1957); Andersson (1971); Ammeter (1971) for fire insurance losses, and Benktander (1962) for automobile insurance losses.

Table 2 provides the KS distances for two parameter sets:  $(\alpha = 2.7163, \beta = 16.8759)$ , and  $(\alpha = 2, \beta = 3000)$  (e.g., Example 5.7 in Bahnemann, 2015). Recall that the  $m^{\text{th}}$  moment of the Pareto distribution is finite only when  $m < \alpha$ . Consequently, most of the existing MMAs are not applicable for these examples; in the first case, only the NA method and MED method with  $m = 2$ , are applicable. In contrast, our approximation is feasible in both cases and works well.

Figure 2 compares the plots of the CDFs in Table 2 against the actual CDF  $F_P$  (in blue). Figures 2 a), and c) depict our approximations (dark green and light green, where light green corresponds to the better approximation) and the MED approximation (in red, when applicable). Figure 2 b) depicts the

normal approximation (orange).

**Example 3.** The final distribution we consider is Weibull, which is a GGC when  $\delta < 1$ , and these are the values of interest when modelling non-life insurance risks (Bahnmann, 2015). Very much like the log-normal and Pareto distributions, the Weibull distribution has been commonly chosen to model insurance data. We refer to Mikosch (2009); Klugman et al. (2012) for a general discussion, as well as to Hogg and Klugman (1983) for an application to hurricane loss data.

Table 3 provides the KS distances for the parameters  $\beta = 220.653$  and  $\delta = 0.8$  (e.g., Example 2.13 in Bahnmann, 2015).

Figure 3 compares the plots of the CDFs in Table 3 with the actual CDF  $F_W$  (in blue). Figure 3 a) depicts our approximations (dark green and light green, where light green corresponds to the better approximation) and the MED approximation (red). Figure 3 b) depicts the simple moment-matching methods: NA (orange), NPA (salmon), and SGA (black).

Table 1: The KS distances,  $d_{KS}(\cdot, X_L)$ , of the approximations to risk RVs distributed log-normally

Parameters	$\tilde{F}_{(5)}$	$\tilde{F}_{(10)}$	$\tilde{F}_{W_{5,70}}$	$\tilde{F}_{NA}$	$\tilde{F}_{NP}$	$\tilde{F}_{SG}$
$\mu = 0, \sigma = 0.5$	9.029e-4	2.950e-6	1.131e-3	9.834e-2	7.165e-2	3.870e-2
Parameters	$\tilde{F}_{(3)}$	$\tilde{F}_{(16)}$	$\tilde{F}_{W_{3,70}}$	$\tilde{F}_{NA}$	$\tilde{F}_{NP}$	$\tilde{F}_{SG}$
$\mu = 1.5240, \sigma = 1.2018$	6.070e-3	2.118e-6	3.400e-2	0.290e0	0.730e0	0.5919e0
Parameters	$\tilde{F}_{(3)}$	$\tilde{F}_{(36)}$	$\tilde{F}_{W_{3,70}}$	$\tilde{F}_{NA}$	$\tilde{F}_{NP}$	$\tilde{F}_{SG}$
$\mu = 5.9809, \sigma = 1.8$	7.968e-3	1.113e-6	-	0.420e0	0.835e0	0.8015e0

Table 2: The KS distances,  $d_{KS}(\cdot, X_P)$ , of the approximations to risk RVs distributed Pareto

Parameters	$\tilde{F}_{(2)}$	$\tilde{F}_{(10)}$	$\tilde{F}_{W_{2,200}}$	$\tilde{F}_{NA}$	$\tilde{F}_{NP}$	$\tilde{F}_{SG}$
$\alpha = 2.7163, \beta = 16.8759$	1.273e-2	4.320e-5	3.477e-2	0.302e0	-	-
Parameters	$\tilde{F}_{(3)}$	$\tilde{F}_{(10)}$	-	$\tilde{F}_{NA}$	$\tilde{F}_{NP}$	$\tilde{F}_{SG}$
$\alpha = 2, \beta = 3000$	6.532e-2	9.525e-3	-	-	-	-

Table 3: The KS distances,  $d_{KS}(\cdot, X_W)$ , of the approximations to a risk RV distributed Weibull


Parameters	$\tilde{F}_{(3)}$	$\tilde{F}_{(10)}$	$\tilde{F}_{W_{3,70}}$	$\tilde{F}_{NA}$	$\tilde{F}_{NP}$	$\tilde{F}_{SG}$
$\beta = 220.653, \delta = 0.8$	3.297e-2	5.337e-4	2.075e-2	2.13e-1	3.45e-1	1.374e-1

Table 4: Computation times for the CDFs in Tables 1, 2, and 3.


Log-normal	$\tilde{F}_{(5)}$	$\tilde{F}_{(10)}$
$\mu = 0, \sigma = 0.5$	19.256	39.120
Log-normal	$\tilde{F}_{(3)}$	$\tilde{F}_{(16)}$
$\mu = 1.5240, \sigma = 1.2018$	14.068	71.052
Log-normal	$\tilde{F}_{(3)}$	$\tilde{F}_{(36)}$
$\mu = 5.9809, \sigma = 1.8$	15.924	162.052
Pareto	$\tilde{F}_{(2)}$	$\tilde{F}_{(10)}$
$\alpha = 2.7163, \beta = 16.8759$	8.876	43.856
Pareto	$\tilde{F}_{(3)}$	$\tilde{F}_{(10)}$
$\alpha = 2, \beta = 3000$	13.308	40.912
Weibull	$\tilde{F}_{(3)}$	$\tilde{F}_{(10)}$
$\beta = 220.653, \delta = 0.8$	12.528	39.852

In summary, we note with satisfaction that the proposed refined MMA method has performed exactly as expected in all of the three examples discussed above. That is, it has outperformed all other MMAs in the cases when the latter are applicable (e.g., the lighter-tailed log-normal CDFs with  $\sigma = 0.5$ ,  $\sigma = 1.2018$  and the Weibull CDF), and it has provided a feasible and accurate alternative, otherwise (e.g., the heavier-tailed log-normal CDF with  $\sigma = 1.8$  and the Pareto CDFs).


We conclude this section by addressing a request of a referee to report the times required to compute the CDFs in Tables 1, 2, and 3 using the MMA approach put forward herein. These times are provided in Table 4. (All calculations were performed on a laptop computer with 12 GB of memory and an Intel® Core™ i5-5200U CPU.) The numbers represent the time, in seconds, required to perform the Gaver Stehfest algorithm described in Section 5 with  $m = 25$  and arithmetic precision to 300 digits and




(a) The CDFs  $F_L$ ,  $\tilde{F}_{(5)}$ ,  $\tilde{F}_{(10)}$ , and  $\tilde{F}_{W_{5,70}}$ .




(b) The CDFs  $F_L$ ,  $\tilde{F}_{NA}$ ,  $\tilde{F}_{NP}$ , and  $\tilde{F}_{SG}$ .




(c) The CDFs  $F_L$ ,  $\tilde{F}_{(3)}$ ,  $\tilde{F}_{(16)}$ , and  $\tilde{F}_{W_{3,70}}$ .



(d) The CDFs  $F_L$ ,  $\tilde{F}_{NA}$ ,  $\tilde{F}_{NP}$ , and  $\tilde{F}_{SG}$ .



(e) The CDFs  $F_L$ ,  $\tilde{F}_{(3)}$ , and  $\tilde{F}_{(36)}$ .




(f) The CDFs  $F_L$ ,  $\tilde{F}_{NA}$ ,  $\tilde{F}_{NP}$ , and  $\tilde{F}_{SG}$ .


Figure 1: The log-normal CDFs and the corresponding approximations as per Table 1.

precomputed coefficients of the approximant Laplace transforms. Remarkably, the computation speed can be significantly enhanced (without dramatic drop in accuracy) by reducing the degree of precision and using a smaller value for  $m$ . Specifically, we set  $m = 10$  and the arithmetic precision to 100 digits.






(a) The CDFs  $F_P$ ,  $\tilde{F}_{(2)}$ ,  $\tilde{F}_{(10)}$ , and  $\tilde{F}_{W_{2,200}}$ .




(b) The CDFs  $F_P$ , and  $\tilde{F}_{NA}$ .




(c) The CDFs  $F_P$ ,  $\tilde{F}_{(3)}$ , and  $\tilde{F}_{(16)}$ .

Figure 2: The Pareto CDFs and the corresponding approximations as per Table 2.



(a) The CDFs  $F_W$ ,  $\tilde{F}_{(3)}$ ,  $\tilde{F}_{(10)}$ , and  $\tilde{F}_{W_{3,70}}$ .



(b) The CDFs  $F_W$ ,  $\tilde{F}_{NA}$ ,  $\tilde{F}_{NP}$ , and  $\tilde{F}_{SG}$ .

Figure 3: The Weibull CDF and the corresponding approximations as per Table 3.

Then in, e.g., the log-normal case with  $\mu = 5.9809$  and  $\sigma = 1.8$ ,  $\tilde{F}_{(3)}$  and  $\tilde{F}_{(36)}$  can be computed in 0.948 and 9.396 seconds, respectively, while achieving the KS distances of  $7.968e-3$  and  $1.123e-6$ , respectively.

We further turn to the aggregate risk RVs, and we demonstrate that the advantages of our approximation method carry on.

## 6.2 Aggregate risks with full and partial coverage

We begin by approximating the CDF of the RV  $S_n = X_1 + \dots + X_n$  - the individual risk model (e.g., [Dhaene and Vincke, 2004](#)), and the CDF of the RV  $S_N = X_1 + \dots + X_N$  - the collective risk model (e.g., [Goovaerts, 2004](#)). In the former case, we sum  $n$  independent RVs with possibly different distributions. In the latter case, we sum a random number  $N$  of IID RVs. Furthermore, as ceding insurers often turn to a reinsurer in order to reduce the variability of the underwriting outcomes, we conclude the discussion in this section with the stop-loss reinsurance set-up. We note briefly that due to [Borch \(1969\)](#), the variance of the cedent insurer's payouts is the smallest under the stop-loss reinsurance contract. Irrespective of whether coverage modifications of the aggregate risks are allowed or not, we assume that the stand-alone risks have CDFs (18), (19), and (20). As explicit CDFs of the aggregate risks are not available, we use the Monte-Carlo (MC) method as a benchmark.

**Example 4.** Consider the CRM,  $S_N = X_1 + \dots + X_N$ , where  $N$  has a Poisson( $\lambda$ ) distribution, and  $X_i$ ,  $i = 1, \dots, N$  has a log-normal distribution with parameters  $\mu = 5.9809$ , and  $\sigma = 1.8$  ([Bahnmann, 2015](#), Example 5.9). The Laplace transform of the RV  $S_N$  is given by

$$\phi_{S_N}(z) = \exp[\lambda(\phi_{X_1}(z) - 1)].$$


Thus we obtain an approximation for  $\phi(z) := \mathbb{E}[\exp(-zS_N)]$  as following

$$\tilde{\phi}(z) = \exp\left[\lambda(\tilde{\phi}_1(z) - 1)\right].$$


Then we evoke the Gaver-Stehfest algorithm to obtain the approximating CDF of the RV  $S_N$ .

Figure 4 summarizes the results. Namely, Figures 4a), c), and e) show the CDFs of our approximation (dark green; succinctly  $\tilde{F}_{S_N, (36)}$ ) and the CDF produced by means of MC simulation with  $10^6$  samples (blue; succinctly  $F_M$ ), for  $\lambda = 5, 10$ , and  $15$ , respectively. In each one of our approximations, we used  $m = 36$  to approximate the underlying log-normal severity distribution. Figures 4 b), d), and f) depict the MC CDF (blue), as well as the CDF obtained via NA (orange), NPA (salmon), and SGA (black).


**Example 5.** Consider the IRM,  $S_n = X_1 + \dots + X_n$ , with  $n = 15$ . We assume  $X_i$ ,  $i = 1, \dots, 5$ , are distributed log-normally with parameters  $\mu = 5.9809, \sigma = 1.8$ ,  $X_i$ ,  $i = 6, \dots, 10$ , are distributed Pareto with parameters  $\alpha = 2, \beta = 3000$ , and  $X_i$ ,  $i = 11, \dots, 15$  are distributed Weibull with parameters  $\beta =$




(a) The CDFs  $\tilde{F}_{S_N,(36)}$  and  $F_M$  with  $\lambda = 5$




(b) The CDFs  $F_M$ ,  $\tilde{F}_{NA}$ ,  $\tilde{F}_{NP}$ , and  $\tilde{F}_{SG}$  with  $\lambda = 5$




(c) The CDFs  $\tilde{F}_{S_N,(36)}$  and  $F_M$  with  $\lambda = 10$



(d) The CDFs  $F_M$ ,  $\tilde{F}_{NA}$ ,  $\tilde{F}_{NP}$ , and  $\tilde{F}_{SG}$  with  $\lambda = 10$ .



(e) The CDFs  $\tilde{F}_{S_N,(36)}$  and  $F_M$  with  $\lambda = 15$



(f) The CDFs  $F_M$ ,  $\tilde{F}_{NA}$ ,  $\tilde{F}_{NP}$ , and  $\tilde{F}_{SG}$  with  $\lambda = 15$

Figure 4: The collective risk model:  $S_N = X_1 + \dots + X_N$ ,  $N \sim \text{Poisson}(\lambda)$ , and the  $X_i$ 's are IID log-normally with the parameters  $(\mu = 5.9809, \sigma = 1.8)$ .

220.653,  $\delta = 0.8$ ; all independent. The LT of the aggregate risk RV  $S_n$ , denoted by  $\phi(z) := \mathbb{E}[\exp(-zS_n)]$ ,

is given by

$$\phi(z) = (\phi_{X_1}(z) \cdot \phi_{X_6}(z) \cdot \phi_{X_{11}}(z))^5.$$

Thus we approximate the CDF of the RV  $S_n$  by first approximating the LTs of the severities and then evoking the Gaver-Stehfest method on

$$\tilde{\phi}(z) = (\tilde{\phi}_1(z) \cdot \tilde{\phi}_6(z) \cdot \tilde{\phi}_{11}(z))^5.$$

Figure 5 shows our approximation of the CDF of  $S_n$  (dark green; denoted by  $\tilde{F}_{S_n,(40),(36),(30)}$ ) and the CDF obtained by means of MC simulation (blue). In this case, we used approximation orders  $m = 40$ ,  $m = 36$ , and  $m = 30$  to approximate the underlying log-normal, Pareto, and Weibull CDFs, respectively. Note that the RV  $S_n$  has only one finite moment and, hence, other MMAs are not applicable.




Figure 5: The CDFs  $\tilde{F}_{S_n,(40),(36),(30)}$  and  $F_M$  for the individual risk model  $S_n$ , with  $n = 15$ .

**Example 6.** In the previous examples we considered aggregate risks with full coverages, that is there were no deductibles, policy limits, or other policy modifications applied by the insurer to reduce the payouts of the benefits. However, this is not always the case. Consequently, in this example, we explore the so-called stop-loss r.v.  $S_{r,l}$ , which is given, for positive retention  $r$  and limit  $l$ , by

$$S_{r,l} = \begin{cases} 0, & X < r \\ S - r, & r \leq X < r + l \\ l, & r + l \leq X < \infty \end{cases}$$

where the RV  $S$  is the aggregate risk RV within either the IRM or CRM. It is a standard exercise to show that the CDF of the RV  $S_{r,l}$  is given by

$$F_{r,l}(s) = \begin{cases} \mathbb{P}[S \leq s + r], & s < l \\ 1, & l \leq s \end{cases}.$$

Consequently, the CDF of the risk RV  $S_{r,l}$  is approximated similarly to the case in Example 4 (if the RV  $S$  refers to the aggregate risk within CRM), and similarly to the case in Example 5 (if the RV  $S$  refers to the aggregate risk within IRM). To demonstrate, we consider the following CRM:  $S := S_N = X_1 + \dots + X_N$ ,  $N \sim \text{Poisson}(15)$ , and the  $X_i$ 's are IID with log-normal distribution and the parameters  $\mu = 5.9809$ ,  $\sigma = 1.8$ . For the sake of the demonstration, we set  $r = \mathbb{E}[N] \cdot 3000 = 45000$ , and  $l = \mathbb{E}[N] \cdot 5000 = 75000$ . Figure 6 depicts the approximating CDF  $\tilde{F}_{r,l}$  of the risk RV  $S_{r,l}$  (dark green) and the CDF obtained by means of the MC simulation  $F_M$  (blue).




Figure 6: The CDFs  $\tilde{F}_{r,l}$  and  $F_M$  for the stop-loss contract with retention  $r = 45000$  and limit  $l = 75000$

## Acknowledgements

We are grateful to the Casualty Actuarial Society (CAS) and the Natural Sciences and Engineering Research Council (NSERC) of Canada for grants supporting our research. Constructive criticism and insightful suggestions of two Referees and the Editor are also greatly appreciated.

## References

- Aase, K. (2004). Pooling in insurance. In Teugels, J. L. and Sundt, B., editors, *Encyclopaedia of Actuarial Science*, pages 1308 – 1311. Wiley, Chichester, England. Vol 3.
- Abate, J. and Whitt, W. (1992). The Fourier-series method for inverting transforms of probability distributions. *Queueing Systems*, 10(1-2):5–87.
- Ammeter, H. (1971). Grosstscha-den-verteilungen und ihre anwendungen, mitt. verein. schweiz. *Versich.-Mathr*, (71):35 – 62.

- Andersson, H. (1971). An analysis of the development of the fire losses in the northern countries after the second world war. *ASTIN Bulletin: the Journal of the International Actuarial Association*, (6):25 – 30.
- Badescu, A., Breuer, L., Da Silva Soares, A., Latouche, G., Remiche, M.-A., and Stanford, D. (2005). Risk processes analyzed as fluid queues. *Scandinavian Actuarial Journal*, 2005(2):127–141.
- Bahnemann, D. (2015). *Distributions for actuaries*. Casualty Actuarial Society.
- Baker, S. G. and Graves-Morris, P. (1996). *Padé Approximants*, volume 1. Cambridge University Press, Cambridge–New York, 2 edition.
- Benckert, L. and Sternberg, I. (1957). An attempt to find an expression for the distribution of fire damage amount. *Transactions of the XV International Congress of Actuaries*, (2):288 – 294.
- Benktander, G. (1962). Notes sur la distribution conditionné du montant d’un sinistre par rapport a l’hypothese qu’il y a eu un sinistre dans l’assurance automobile. *ASTIN Bulletin: the Journal of the International Actuarial Association*, (2):24 – 29.
- Bickerstaff, D. R. (1972). Automobile collision deductibles and repair cost groups: the log-normal model. *Proceedings of the Casualty Actuarial Society*, LIX:68.
- Bondesson, L. (1992). *Generalized Gamma Convolutions and Related Classes of Distributions and Densities*, volume 76 of *Lecture Notes in Statistics*. Springer.
- Borch, K. (1969). The optimal reinsurance treaty. *ASTIN Bulletin: the Journal of the International Actuarial Association*, (5):293 – 297.
- Brockett, P. (1983). On the misuse of the central limit theorem in some risk calculations. *Journal of Risk and Insurance*, 50(4):727 – 731.
- Cossette, H., Landriault, D., Marceau, E., and Moutanabbir, K. (2016). Moment-based approximation with mixed Erlang distributions. *Variance*, 10(1):161–182.
- Davies, B. and Martin, B. (1979). Numerical inversion of the Laplace transform: a survey and comparison of methods. *Journal of Computational Physics*, 33(1):1 – 32.
- Dhaene, J. and Vincke, D. (2004). Individual risk models. In Teugels, J. L. and Sundt, B., editors, *Encyclopaedia of Actuarial Science*, pages 871 – 875. Wiley, Chichester, England. Vol 2.
- Dropkin, L. B. (1964). Size of loss distributions in workmens compensation insurance. *Proceedings of the Casualty Actuarial Society*, LI:198.

- Dufresne, D. (2007). Fitting combinations of exponential to probability distributions,. *Applied Stochastic Models in Business and Industry*, (23):23 – 48.
- EIOPA-14-322 (2014). The underlying assumptions in the standard formula for the Solvency Capital Requirement calculation. Technical report, European Insurance and Occupational Pensions Authority.
- Furman, E., Hackmann, D., and Kuznetsov, A. (2018). On log-normal convolutions: An analytical-numerical method with applications to economic capital determination. *Insurance: Mathematics and Economics*. under review.
- Goovaerts, M. (2004). Collective risk models. In Teugels, J. L. and Sundt, B., editors, *Encyclopaedia of Actuarial Science*. Wiley, Chichester, England.
- Hardy, M. R. (2004). Approximating the aggregate claims distributions. In Teugels, J. L. and Sundt, B., editors, *Encyclopaedia of Actuarial Science*, pages 78 – 86. Wiley, Chichester, England. Vol 1.
- Hogg, R. and Klugman, S. (1983). On the estimation of long tailed skewed distributions with actuarial applications. *Journal of Econometrics*, (23):91 – 102.
- Kleiber, C. and Kotz, S. (2003). *Statistical Size Distributions in Economics and Actuarial Sciences*. John Wiley & Sons, Inc., Hoboken, New Jersey. Canada.
- Klugman, S. A., Panjer, H. H., and Willmot, G. E. (2012). *Loss models: from data to decisions*, 3rd ed. Society of Actuaries.
- Kou, S. G. and Wang, H. (2003). First passage times of a jump diffusion process. *Advances in Applied Probability*, 35(2):pp. 504–531.
- Kuznetsov, A. (2013). On the convergence of the gaver-stehfest algorithm. *SIAM Journal on Numerical Analysis*, 6(51):2984 – 2998.
- Mikosch, T. (2009). *Non-Life Insurance Mathematics: An Introduction with Poisson Process*, volume 32. SIAM Review.
- O’Neill, B. and Wells, W. T. (1972). Some Recent Results in Lognormal Parameter Estimation Using Grouped and Ungrouped Data. *Journal of the American Statistical Association*, 67(337):76.
- Ramsay, C. (1991). A note of the normal power approximation. *ASTIN Bulletin: the Journal of the International Actuarial Association*, (21):147 – 150.
- Schoutens, W. and Damme, G. V. (2011). The  $\beta$ -variance gamma model. *Review of Derivatives Research*, 14(3):263–282.

- Seal, H. (1980). Survival probabilities based on pareto claim distributions. *ASTIN Bulletin: the Journal of the International Actuarial Association*, (11):61 – 71.
- Stehfest, H. (1970). Algorithm 368: Numerical inversion of Laplace transforms. *Commun. ACM*, 13(1):47–49.
- Takahasi, H. and Mori, M. (1974). Double exponential formulas for numerical integration. *Publ. RIMS, Kyoto Univ.*, 9:721–741.
- Tao, J., Provost, S., and Ren, J. (2016). Moment-based density approximations for aggregate losses. *Scandinavian Actuarial Journal*, 2016(3):216 – 245.
- Thorin, O. (1977). On the infinite divisibility of the lognormal distribution. *Scandinavian Actuarial Journal*, 1977(3):121–148.